

Advanced Quantum Mechanics

Prof. Dr. Stefan Dittmaier

Albert-Ludwigs-Universität Freiburg, Physikalisches Institut
D-79104 Freiburg, Germany

Winter-Semester 2018/19

Draft: February 4, 2019

Contents

1	Recapitulation of basic qm. principles	5
2	Symmetries in quantum mechanics	7
3	Approximation methods	9
4	Scattering theory	11
5	Quantization of the electromagnetic field	13
5.1	Free electromagnetic fields	14
5.1.1	Classical description	14
5.1.2	Quantization	17
5.2	Interacting electromagnetic fields	22
5.2.1	Classical fields	22
5.2.2	Quantization	23
5.2.3	Application: $1e^-$ atoms in quantized radiation field	25

Chapter 5

Quantization of the electromagnetic field

Experimental observation:

Elmg. fields of frequency ν emits / absorbs *energy* in portions $h\nu = \hbar\omega$.

Elmg. fields of frequency ν emits / absorbs *momentum* in portions $h/\lambda = \hbar k$.

⇒ Qm. principles should be applied to the elmg. fields !

5.1 Free electromagnetic fields

5.1.1 Classical description

Maxwell's equations:

$$\begin{aligned}\vec{\nabla} \cdot \vec{E} &= \rho/\epsilon_0, & \vec{\nabla} \times \vec{E} &= -\dot{\vec{B}}, \\ \vec{\nabla} \cdot \vec{B} &= 0, & \vec{\nabla} \times \vec{B} &= \dot{\vec{E}}/c^2 + \mu_0 \vec{j},\end{aligned}\quad (5.1)$$

where the electric charge ρ and the current density \vec{j} vanish for free fields: $\rho = 0, \vec{j} = 0$.

Gauge potentials \vec{A} and Φ :

↪ eliminate the homogenous Maxwell eqs. by construction:

$$\vec{B} = \vec{\nabla} \times \vec{A}, \quad \vec{E} = -\vec{\nabla} \Phi - \dot{\vec{A}}. \quad (5.2)$$

Field eqs. for \vec{A} and Φ :

$$\square \vec{A} + \vec{\nabla}(\vec{\nabla} \cdot \vec{A} + \dot{\Phi}/c^2) = \mu_0 \vec{j}, \quad \triangle \Phi + \vec{\nabla} \cdot \dot{\vec{A}} = -\rho/\epsilon_0. \quad (5.3)$$

Recall: $\square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \triangle$ = wave operator

Elmg. gauge invariance:

Field strengths \vec{E} and \vec{B} (which define the physical state of the classial system) are invariant under the “gauge transformation”:

$$\vec{A} \rightarrow \vec{A}' = \vec{A} + \vec{\nabla} \chi, \quad \Phi \rightarrow \Phi' = \Phi - \dot{\chi}, \quad (5.4)$$

with χ denoting any function $\chi = \chi(\vec{x}, t)$.

⇒ Field eqs. can be simplified by “fixing a gauge”, e.g.:

- $\vec{\nabla} \cdot \vec{A} = 0$, “Coulomb gauge” (used in the following),
- $\vec{\nabla} \cdot \vec{A} + \dot{\Phi}/c^2 = 0$, “Lorenz gauge” (appropriate in relativistic theories).

Free elmg. radiation in Coulomb gauge (“radiation gauge”):

$$\Phi \equiv 0, \quad (5.5)$$

$$\square \vec{A} = 0, \quad \text{homogenous wave equation}, \quad (5.6)$$

$$\vec{\nabla} \vec{A} = 0. \quad (5.7)$$

Basis solution for finite volume V (cubic box of side length L):

- Ansatz: $\vec{A}(\vec{x}, t) = \vec{\varepsilon} e^{i\vec{k}\vec{x} - i\omega t}$, $\vec{\varepsilon} = \text{const.}$
- From Eq.(5.6): $-\frac{\omega^2}{c^2} + \vec{k}^2 = 0$, i.e. $\omega = c|\vec{k}| = ck$.
- From periodicity: $\vec{k} = \frac{2\pi}{L}\vec{n}$, $n_j \in \mathbb{Z}$, i.e. all \vec{k} on discrete lattice.
- From Eq.(5.7): $\vec{k} \cdot \vec{\varepsilon} = 0 \rightarrow 2$ independent solutions (“polarizations”) for each \vec{k} . Convenient choice of $\vec{\varepsilon}_\lambda(\vec{k})$: “helicity basis” $\vec{\varepsilon}_\pm(\vec{k})$.

$$\vec{\varepsilon}_\pm = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ \pm i \\ 0 \end{pmatrix} \quad \text{for } \vec{k} = k\vec{e}_3, \quad \text{other directions via rotations}, \quad (5.8)$$

i.e. $\vec{\varepsilon}_\pm$ describe right/left-circular polarization,

$$\text{normalization: } \vec{\varepsilon}_\lambda \cdot \vec{\varepsilon}_{\lambda'}^* = \delta_{\lambda\lambda'}, \quad \vec{\varepsilon}_\pm^* = \vec{\varepsilon}_\mp, \quad (5.9)$$

$$\text{completeness relation: } \sum_{\lambda=\pm} \varepsilon_{\lambda,a}(\vec{k}) \varepsilon_{\lambda,b}^*(\vec{k}) = \delta_{ab} - \frac{k_a k_b}{k^2}. \quad (5.10)$$

\Rightarrow General solution for $\vec{A}(\vec{x}, t)$:

$$\vec{A}(\vec{x}, t) = \sum_{\vec{k}} \sum_{\lambda} \frac{1}{\sqrt{2\omega\epsilon_0 V}} \left(\underbrace{a_\lambda(\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t}}_{\text{arbitrary amplitudes}} + \underbrace{\text{c.c.}}_{\text{complex conjugate}} \right). \quad (5.11)$$

(normalization defined with some foresight)

Field strengths derived from $\vec{A}(\vec{x}, t)$:

$$\vec{E}(\vec{x}, t) = i \sum_{\vec{k}} \sum_{\lambda} \sqrt{\frac{\omega}{2\epsilon_0 V}} \left(a_\lambda(\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} - \text{c.c.} \right), \quad (5.12)$$

$$\vec{B}(\vec{x}, t) = i \sum_{\vec{k}} \sum_{\lambda} \frac{1}{\sqrt{2\omega\epsilon_0 V}} \left(a_\lambda(\vec{k}) \vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} - \text{c.c.} \right). \quad (5.13)$$

Energy H_{rad} and momentum \vec{P}_{rad} of the field configuration:

$$\begin{aligned}
 H_{\text{rad}} &= \int_V d^3x \frac{1}{2} \left(\epsilon_0 \vec{E}^2 + \vec{B}^2 / \mu_0 \right) \\
 &= -\frac{1}{2V} \int_V d^3x \sum_{\vec{k}, \vec{k}', \lambda, \lambda'} \left\{ \frac{\sqrt{\omega\omega'}}{2} \left(a_\lambda(\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x}-i\omega t} - \text{c.c.} \right) \cdot \left(a_{\lambda'}(\vec{k}') \vec{\varepsilon}_{\lambda'}(\vec{k}') e^{i\vec{k}'\vec{x}-i\omega' t} - \text{c.c.} \right) \right. \\
 &\quad \left. + \underbrace{\frac{1}{2\epsilon_0\mu_0\sqrt{\omega\omega'}}}_{=1/c^2} \left(a_\lambda(\vec{k}) \vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x}-i\omega t} - \text{c.c.} \right) \cdot \left(a_{\lambda'}(\vec{k}') \vec{k}' \times \vec{\varepsilon}_{\lambda'}(\vec{k}') e^{i\vec{k}'\vec{x}-i\omega' t} - \text{c.c.} \right) \right\} \\
 &= -\frac{1}{4V} \int_V d^3x \sum_{\vec{k}, \vec{k}', \lambda, \lambda'} \left\{ \sqrt{\omega\omega'} \left(a_\lambda(\vec{k}) a_{\lambda'}(\vec{k}') \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}(\vec{k}') \underbrace{e^{i(\vec{k}+\vec{k}')\vec{x}}}_{\int d^3x \rightarrow V\delta_{\vec{k},-\vec{k}'} e^{-i(\omega+\omega')t}} \right. \right. \\
 &\quad \left. \left. - a_\lambda(\vec{k}) a_{\lambda'}^*(\vec{k}') \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}^*(\vec{k}') e^{i(\vec{k}-\vec{k}')\vec{x}-i(\omega-\omega')t} + \text{c.c.} \right) \right. \\
 &\quad \left. + \frac{c^2}{\sqrt{\omega\omega'}} \left[a_\lambda(\vec{k}) a_{\lambda'}(\vec{k}') \left(\vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) \right) \cdot \left(\vec{k}' \times \vec{\varepsilon}_{\lambda'}(\vec{k}') \right) e^{i(\vec{k}+\vec{k}')\vec{x}-i(\omega+\omega')t} \right. \right. \\
 &\quad \left. \left. - a_\lambda(\vec{k}) a_{\lambda'}^*(\vec{k}') \left(\vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) \right) \cdot \left(\vec{k}' \times \vec{\varepsilon}_{\lambda'}^*(\vec{k}') \right) e^{i(\vec{k}-\vec{k}')\vec{x}-i(\omega-\omega')t} + \text{c.c.} \right] \right\} \\
 &= -\frac{1}{4} \sum_{\vec{k}, \lambda, \lambda'} \left\{ \omega \left(\underbrace{a_\lambda(\vec{k}) a_{\lambda'}(-\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}(-\vec{k}) e^{-2i\omega t}}_{=k^2 \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}(-\vec{k})} - a_\lambda(\vec{k}) a_{\lambda'}^*(\vec{k}) \underbrace{\vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}^*(\vec{k})}_{=\delta_{\lambda\lambda'}} + \text{c.c.} \right) \right. \\
 &\quad \left. + \frac{c^2}{\omega} \left[\underbrace{a_\lambda(\vec{k}) a_{\lambda'}(-\vec{k}) \left(\vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) \right) \cdot \left(-\vec{k} \times \vec{\varepsilon}_{\lambda'}(-\vec{k}) \right) e^{-2i\omega t}}_{=-k^2 \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}(-\vec{k})} \right. \right. \\
 &\quad \left. \left. - a_\lambda(\vec{k}) a_{\lambda'}^*(\vec{k}) \underbrace{\left(\vec{k} \times \vec{\varepsilon}_\lambda(\vec{k}) \right) \cdot \left(\vec{k} \times \vec{\varepsilon}_{\lambda'}^*(\vec{k}) \right)}_{=k^2 \vec{\varepsilon}_\lambda(\vec{k}) \cdot \vec{\varepsilon}_{\lambda'}^*(\vec{k}) = \frac{\omega^2}{c^2} \delta_{\lambda\lambda'}} + \text{c.c.} \right] \right\} \\
 &= \sum_{\vec{k}, \lambda} \frac{\omega}{2} \left(a_\lambda(\vec{k}) a_\lambda(\vec{k})^* + a_\lambda(\vec{k})^* a_\lambda(\vec{k}) \right), \tag{5.14}
 \end{aligned}$$

$$\vec{P}_{\text{rad}} = \int_V d^3x \epsilon_0 \vec{E} \times \vec{B} = \dots = \sum_{\vec{k}, \lambda} \frac{\vec{k}}{2} \left(a_\lambda(\vec{k}) a_\lambda(\vec{k})^* + a_\lambda(\vec{k})^* a_\lambda(\vec{k}) \right). \tag{5.15}$$

5.1.2 Quantization

Comparison with system of harmonic oscillators:

$$\hat{H} = \sum_k \left(\frac{\hat{p}_k^2}{2m_k} + \frac{m_k\omega_k^2}{2} \hat{q}_k^2 \right), \quad [\hat{q}_k, \hat{p}_{k'}] = i\hbar\delta_{kk'}, \quad (5.16)$$

$$= \sum_k \frac{\hbar\omega_k}{2} \left(\hat{a}_k \hat{a}_k^\dagger + \hat{a}_k^\dagger \hat{a}_k \right), \quad [\hat{a}_k, \hat{a}_{k'}^\dagger] = \delta_{kk'}, \quad [\hat{a}_k, \hat{a}_{k'}] = [\hat{a}_k^\dagger, \hat{a}_{k'}^\dagger] = 0, \quad (5.17)$$

$$E_{n_1 n_2 \dots} = \sum_k \hbar\omega_k \left(n_k + \frac{1}{2} \right), \quad n_k = 0, 1, 2, \dots \quad (5.18)$$

\Rightarrow Promote classical amplitudes $a_\lambda(\vec{k})$ and $a_\lambda(\vec{k})^*$ to annihilation and creation operators $\sqrt{\hbar} \hat{a}_\lambda(\vec{k})$ and $\sqrt{\hbar} \hat{a}_\lambda(\vec{k})^\dagger$:

- Commutators:

$$[\hat{a}_\lambda(\vec{k}), \hat{a}_{\lambda'}(\vec{k}')^\dagger] = \delta_{\lambda\lambda'} \delta_{\vec{k}\vec{k}'}, \quad [\hat{a}_\lambda(\vec{k}), \hat{a}_{\lambda'}(\vec{k}')] = [\hat{a}_\lambda(\vec{k})^\dagger, \hat{a}_{\lambda'}(\vec{k}')^\dagger] = 0. \quad (5.19)$$

- Hamilton operator:

$$\hat{H}_{\text{rad}} = \sum_{\vec{k}, \lambda} \frac{\hbar\omega}{2} \left(\hat{a}_\lambda(\vec{k}) \hat{a}_\lambda(\vec{k})^\dagger + \hat{a}_\lambda(\vec{k})^\dagger \hat{a}_\lambda(\vec{k}) \right). \quad (5.20)$$

- Field operator:

$$\begin{aligned} \hat{\vec{A}}(\vec{x}, t) &= \sum_{\vec{k}} \sum_{\lambda} \sqrt{\frac{\hbar}{2\omega\epsilon_0 V}} \left(\hat{a}_\lambda(\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} + \hat{a}_\lambda(\vec{k})^\dagger \vec{\varepsilon}_\lambda^*(\vec{k}) e^{-i\vec{k}\vec{x} + i\omega t} \right) \\ &= \hat{\vec{A}}(\vec{x}, t)^\dagger = \text{hermitian}. \end{aligned} \quad (5.21)$$

Perform continuum limit $V \rightarrow \infty$:

$$\begin{aligned} \delta_{\vec{k}\vec{k}'} &\rightarrow \frac{(2\pi)^3}{V} \delta(\vec{k} - \vec{k}'), \\ \sum_{\vec{k}} &\rightarrow V \int \frac{d^3 k}{(2\pi)^3}, \\ \text{rescaling: } a_\lambda(\vec{k})^{(\dagger)} &\rightarrow a_\lambda(\vec{k})^{(\dagger)} / \sqrt{V}. \end{aligned} \quad (5.22)$$

⇒ Results:

- Commutators:

$$[\hat{a}_\lambda(\vec{k}), \hat{a}_{\lambda'}(\vec{k}')^\dagger] = (2\pi)^3 \delta_{\lambda\lambda'} \delta(\vec{k} - \vec{k}'), \quad [\hat{a}_\lambda(\vec{k}), \hat{a}_{\lambda'}(\vec{k}')] = [\hat{a}_\lambda(\vec{k})^\dagger, \hat{a}_{\lambda'}(\vec{k}')^\dagger] = 0. \quad (5.23)$$

- Field operators:

$$\hat{\vec{A}}(\vec{x}, t) = \int \frac{d^3k}{(2\pi)^3} \sum_\lambda \sqrt{\frac{\hbar}{2\omega\epsilon_0}} \left(\hat{a}_\lambda(\vec{k}) \vec{\varepsilon}_\lambda(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} + \hat{a}_\lambda(\vec{k})^\dagger \vec{\varepsilon}_\lambda^*(\vec{k}) e^{-i\vec{k}\vec{x} + i\omega t} \right), \quad (5.24)$$

$$\vec{\nabla} \hat{\vec{A}}(\vec{x}, t) = 0, \quad \text{Coulomb gauge condition,} \quad (5.25)$$

$$\hat{\vec{E}}(\vec{x}, t) = -\frac{\partial \hat{\vec{A}}}{\partial t}(\vec{x}, t), \quad (5.26)$$

$$\hat{\vec{B}}(\vec{x}, t) = \vec{\nabla} \times \hat{\vec{A}}(\vec{x}, t). \quad (5.27)$$

- Hamilton operator:

$$\hat{H}_{\text{rad}} = \int \frac{d^3k}{(2\pi)^3} \sum_\lambda \frac{\hbar\omega}{2} \left(\hat{a}_\lambda(\vec{k}) \hat{a}_\lambda(\vec{k})^\dagger + \hat{a}_\lambda(\vec{k})^\dagger \hat{a}_\lambda(\vec{k}) \right). \quad (5.28)$$

- Operator for field momentum:

$$\hat{\vec{P}}_{\text{rad}} = \int \frac{d^3k}{(2\pi)^3} \sum_\lambda \frac{\hbar\vec{k}}{2} \left(\hat{a}_\lambda(\vec{k}) \hat{a}_\lambda(\vec{k})^\dagger + \hat{a}_\lambda(\vec{k})^\dagger \hat{a}_\lambda(\vec{k}) \right). \quad (5.29)$$

Considerations about the field operators:

- By construction, the field operators obey their “eqs. of motion” (EOM) and thus are defined in the Heisenberg picture of time evolution.
- The quantized elmg. field defines a qm. system with infinitely many degrees of freedom:
 - The field modes characterized by \vec{k} and λ are independent harmonic oscillators.
 - Alternatively, the field can be interpreted to allow for excitations at each point \vec{x} , but the fields at different points are NOT independent (derivatives in EOM correspond to interactions of neighbouring points).

- Still to be clarified:

What is the canonical momentum variable corresponding to $\hat{A}(\vec{x}, t)$?

→ Use again analogy to harmonic oscillator:

$$\hat{q}_k = \sqrt{\frac{\hbar}{2m_k\omega_k}} (\hat{a}_k + \hat{a}_k^\dagger) \longrightarrow \hat{A}(\vec{x}, t),$$

$$\hat{p}_k = i\sqrt{\frac{\hbar m_k \omega_k}{2}} (\hat{a}_k^\dagger - \hat{a}_k) \longrightarrow \frac{\partial \hat{A}}{\partial t}(\vec{x}, t) ? \quad (\text{because of factor "i" and sign change})$$

Calculation of commutator (at equal times!):

$$\begin{aligned} \left[\hat{A}_a(\vec{x}, t), \frac{\partial \hat{A}_b}{\partial t}(\vec{y}, t) \right] &= \int \frac{d^3 k}{(2\pi)^3} \int \frac{d^3 k'}{(2\pi)^3} \frac{i\omega' \hbar}{2\epsilon_0 \sqrt{\omega \omega'}} \sum_{\lambda, \lambda'} \\ &\times \left[\left(\hat{a}_\lambda(\vec{k}) \varepsilon_{\lambda, a}(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} + \hat{a}_\lambda(\vec{k})^\dagger \varepsilon_{\lambda, a}^*(\vec{k}) e^{-i\vec{k}\vec{x} + i\omega t} \right), \right. \\ &\quad \left. \left(-\hat{a}_{\lambda'}(\vec{k}') \varepsilon_{\lambda', b}(\vec{k}') e^{i\vec{k}'\vec{y} - i\omega' t} + \hat{a}_{\lambda'}(\vec{k}')^\dagger \varepsilon_{\lambda', b}^*(\vec{k}') e^{-i\vec{k}'\vec{y} + i\omega' t} \right) \right] \\ &= \int \frac{d^3 k}{(2\pi)^3} \int \frac{d^3 k'}{(2\pi)^3} \frac{i\omega' \hbar}{2\epsilon_0 \sqrt{\omega \omega'}} \sum_{\lambda, \lambda'} \\ &\times \left(\underbrace{\left[\hat{a}_\lambda(\vec{k}), \hat{a}_{\lambda'}(\vec{k}')^\dagger \right]}_{=(2\pi)^3 \delta_{\lambda\lambda'} \delta(\vec{k} - \vec{k}')} \varepsilon_{\lambda, a}(\vec{k}) \varepsilon_{\lambda', b}^*(\vec{k}') e^{i\vec{k}\vec{x} - i\vec{k}'\vec{y} - i(\omega - \omega')t} \right. \\ &\quad \left. - \left[\hat{a}_\lambda(\vec{k})^\dagger, \hat{a}_{\lambda'}(\vec{k}') \right] \varepsilon_{\lambda, a}^*(\vec{k}) \varepsilon_{\lambda', b}(\vec{k}') e^{-i\vec{k}\vec{x} + i\vec{k}'\vec{y} + i(\omega - \omega')t} \right) \\ &= \frac{i\hbar}{2\epsilon_0} \int \frac{d^3 k}{(2\pi)^3} \left(\underbrace{\sum_\lambda \varepsilon_{\lambda, a}(\vec{k}) \varepsilon_{\lambda, b}^*(\vec{k}) e^{i\vec{k}(\vec{x} - \vec{y})}}_{=\delta_{ab} - \frac{k_a k_b}{k^2}} + \text{c.c.} \right) \\ &= \frac{i\hbar}{\epsilon_0} \int \frac{d^3 k}{(2\pi)^3} \left(\underbrace{\delta_{ab} - \frac{k_a k_b}{k^2}}_{\equiv \delta_{ab}^\perp(\vec{x} - \vec{y}), \text{ "transverse } \delta\text{-function}} \right) e^{i\vec{k}(\vec{x} - \vec{y})}. \end{aligned}$$

⇒ Identification of conjugate momentum variable: $\hat{\vec{\Pi}}(\vec{x}, t) \equiv \epsilon_0 \frac{\partial \hat{A}}{\partial t}(\vec{x}, t) = -\epsilon_0 \hat{\vec{E}}(\vec{x}, t)$.

$$\left[\hat{A}_a(\vec{x}, t), \hat{\Pi}_b(\vec{y}, t) \right] = i\hbar \delta_{ab}^\perp(\vec{x} - \vec{y}). \quad (5.30)$$

Comments:

- Definition of $\vec{\Pi}(\vec{x}, t)$ should be verified upon checking the canonical EOMs.
- $\delta_{ab}^\perp(\vec{x}) = 3 \times 3$ matrix-valued projector on transverse vector fields:

$$\delta_{ab}^\perp(\vec{x}) = \int \frac{d^3k}{(2\pi)^3} \left(\delta_{ab} - \frac{k_a k_b}{k^2} \right) e^{ik\vec{x}} = \delta_{ab}\delta(\vec{x}) + \frac{\partial^2}{\partial x_a \partial x_b} \frac{1}{4\pi|\vec{x}|}, \quad (5.31)$$

$$V_a^\perp(\vec{x}) \equiv \int d^3y \sum_b \delta_{ab}^\perp(\vec{x} - \vec{y}) V_b(\vec{y}) = V_a(\vec{x}) + \int d^3y \frac{\partial}{\partial x_a} \frac{\vec{\nabla}_y \vec{V}(\vec{y})}{4\pi|\vec{x} - \vec{y}|}.$$

$$\hookrightarrow \vec{\nabla} \vec{V}^\perp(\vec{x}) = 0. \quad (5.32)$$

$$\sum_a \frac{\partial}{\partial x_a} \delta_{ab}^\perp(\vec{x}) = 0, \quad \text{Tr} \{ \delta^\perp(\vec{x}) \} = 2\delta(\vec{x}). \quad (5.33)$$

- Relativistic covariance of quantization in Coulomb gauge maintained, but non-trivial to prove.
- Manifestly relativistically covariant quantization possible (“Gupta–Bleuler method”):
 - ◊ $[\hat{A}^\mu(\vec{x}, t), \hat{\Pi}^\nu(\vec{y}, t)] = i\hbar g^{\mu\nu} \delta(\vec{x} - \vec{y})$ for field operators.,
 - ◊ $\langle \phi | \partial_\mu \hat{A}^\mu | \phi \rangle = 0$ for states $|\phi\rangle$.

\hookrightarrow Lecture on relativistic QFT !

“Fock space” of photons (follows analogy to harmonic oscillator)

- “vacuum state” $|0\rangle$ (no photons): $\langle 0|0\rangle = 1$, $a_\lambda(\vec{k})|0\rangle = 0 \quad \forall \lambda, \vec{k}$.

$$\Rightarrow E_{\text{vac}} = \langle 0|\hat{H}_{\text{rad}}|0\rangle = \sum_{\vec{k}, \lambda} \frac{\hbar\omega}{2} \underbrace{\langle 0|\hat{a}_\lambda(\vec{k})\hat{a}_\lambda(\vec{k})^\dagger + \hat{a}_\lambda(\vec{k})^\dagger\hat{a}_\lambda(\vec{k})|0\rangle}_{=1} = \sum_{\vec{k}, \lambda} \frac{\hbar\omega}{2} \rightarrow \infty,$$

$$\langle 0|\hat{\vec{P}}_{\text{rad}}|0\rangle = \sum_{\vec{k}, \lambda} \frac{\hbar\vec{k}}{2} = 0. \quad (5.34)$$

Note: Meaning of infinite vacuum energy not really fully understood,
but not very problematic in practice, since not directly measurable.

Measured: “Casimir effect” = force between two electrically neutral metal plates
= change of vacuum energy with changing volume.

↪ Redefine \hat{H}_{rad} by splitting off unobservable E_{vac} :

$$\hat{H}_{\text{rad}} \rightarrow \hat{H}_{\text{rad}} - E_{\text{vac}}. \quad \Rightarrow \hat{H}_{\text{rad}} = \sum_{\vec{k}, \lambda} \hbar\omega \hat{a}_\lambda(\vec{k})^\dagger \hat{a}_\lambda(\vec{k}). \quad (5.35)$$

- 1-photon states: $|\vec{k}, \lambda\rangle \equiv \hat{a}_\lambda(\vec{k})^\dagger|0\rangle$, $\langle \vec{k}, \lambda | \vec{k}', \lambda' \rangle = (2\pi)^3 \delta_{\lambda\lambda'} \delta(\vec{k} - \vec{k}')$.

$$\begin{aligned} \hat{H}_{\text{rad}}|\vec{k}, \lambda\rangle &= \sum_{\vec{k}', \lambda'} \hbar\omega' \hat{a}_{\lambda'}(\vec{k}')^\dagger \hat{a}_{\lambda'}(\vec{k}') \hat{a}_\lambda(\vec{k})^\dagger|0\rangle \\ &= \sum_{\vec{k}', \lambda'} \hbar\omega' \left(\hat{a}_{\lambda'}(\vec{k}')^\dagger \underbrace{\left[\hat{a}_{\lambda'}(\vec{k}'), \hat{a}_\lambda(\vec{k})^\dagger \right]}_{=\delta_{\lambda\lambda'}\delta_{\vec{k}\vec{k}'}} + \overbrace{\hat{a}_{\lambda'}(\vec{k}')^\dagger \hat{a}_\lambda(\vec{k})^\dagger|0\rangle \hat{a}_{\lambda'}(\vec{k}')} \right) |0\rangle \\ &= \hbar\omega \hat{a}_\lambda(\vec{k})^\dagger|0\rangle = \hbar\omega |\vec{k}, \lambda\rangle, \end{aligned}$$

$$\hat{\vec{P}}_{\text{rad}}|\vec{k}, \lambda\rangle = \hbar\vec{k}|\vec{k}, \lambda\rangle. \quad (5.36)$$

Polarization clarified upon investigating behaviour of $|\vec{k}, \lambda\rangle$ under rotations.

↪ $|\vec{k}, \lambda\rangle$ = state with 1 photon of energy $\hbar\omega$, momentum $\hbar\vec{k}$, and polarization $\vec{\varepsilon}_\lambda(\vec{k})$.

- N -photon state:

$$|\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle \propto \left(\prod_{n=1}^N \hat{a}_{\lambda_n}(\vec{k}_n)^\dagger \right) |0\rangle. \quad (5.37)$$

$|\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle$ = symmetric under exchange of any pair (\vec{k}_i, λ_i) and (\vec{k}_j, λ_j) .

⇒ Bosonic states !

$$\begin{aligned} \hat{H}_{\text{rad}}|\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle &= \sum_{n=1}^N \hbar\omega_n |\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle, \\ \hat{\vec{P}}_{\text{rad}}|\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle &= \sum_{n=1}^N \hbar\vec{k}_n |\vec{k}_1, \lambda_1; \dots; \vec{k}_N, \lambda_N\rangle. \end{aligned} \quad (5.38)$$

“Fock space” \mathcal{F} = Hilbert space spanned by all multi-photon states.

5.2 Interacting electromagnetic fields

5.2.1 Classical fields

Considered system: N particles with masses m_n and electric charges q_n at positions \vec{x}_n ($n = 1, 2, \dots, N$).

Charge density: $\rho(\vec{x}) = \sum_n q_n \delta(\vec{x} - \vec{x}_n)$.

Electric field \vec{E} and potentials in Coulomb gauge ($\vec{\nabla} \vec{A} = 0$):

$$\vec{E} = \underbrace{-\vec{\nabla}\Phi}_{=\vec{E}^{\parallel}} - \underbrace{\dot{\vec{A}}}_{=\vec{E}^{\perp}} = \vec{E}^{\parallel} + \vec{E}^{\perp}. \quad (5.39)$$

$$\vec{\nabla} \vec{E} = \vec{\nabla} \vec{E}^{\parallel} = -\Delta \Phi = \rho/\epsilon_0. \quad \rightarrow \quad \Phi(\vec{x}) = \sum_n \frac{q_n}{4\pi\epsilon_0 |\vec{x} - \vec{x}_n|} = \text{no dynamical variable!}$$

$$\vec{E}^{\perp} = -\dot{\vec{A}}, \quad \vec{\nabla} \vec{E}^{\perp} = 0, \quad \vec{\Pi} = \epsilon_0 \dot{\vec{A}} = -\epsilon_0 \vec{E}^{\perp} = \text{canonical momentum variable to } \vec{A}.$$

Hamilton function:

$$H = H_{\text{matter}} + H_{\text{elmg}}, \quad (5.40)$$

$$H_{\text{matter}}(\{\vec{x}_n, \vec{p}_n\}) = \sum_n \frac{1}{2m_n} \left(\underbrace{\vec{p}_n - q_n \vec{A}(\vec{x}_n, t)}_{= m_n \dot{\vec{x}}_n} \right)^2, \quad (5.41)$$

$$\begin{aligned} H_{\text{elmg}}(\vec{A}, \vec{\Pi}) &= \int_V d^3x \frac{1}{2} \left(\epsilon_0 \vec{E}^{\parallel 2} + \vec{B}^2 / \mu_0 \right) \\ &= \int_V d^3x \frac{1}{2} \left(\epsilon_0 (\vec{E}^{\perp})^2 + 2\epsilon_0 \underbrace{\vec{E}^{\perp} \vec{E}^{\parallel}}_{\substack{\equiv (\vec{\nabla} \vec{E}^{\perp}) \Phi = 0 \\ \text{P.I.}}} + \underbrace{\epsilon_0 (\vec{E}^{\parallel})^2}_{\substack{\equiv -\epsilon_0 \Phi \Delta \Phi = \rho \Phi}} + \vec{B}^2 / \mu_0 \right) \\ &= \int_V d^3x \frac{1}{2} \left(\epsilon_0 (\vec{E}^{\perp})^2 + \vec{B}^2 / \mu_0 \right) + \frac{1}{2} \sum_n q_n \Phi(\vec{x}_n) \\ &= \underbrace{\int_V d^3x \frac{1}{2} \left(\vec{\Pi}^2 / \epsilon_0 + (\vec{\nabla} \times \vec{A})^2 / \mu_0 \right)}_{\equiv H_{\text{rad}}(\vec{A}, \vec{\Pi})} + V_{\text{Coul}}(\{\vec{x}_n\}), \end{aligned} \quad (5.42)$$

$$V_{\text{Coul}}(\{\vec{x}_n\}) = \frac{1}{2} \sum_{m \neq n} \frac{q_m q_n}{4\pi\epsilon_0 |\vec{x}_m - \vec{x}_n|} + \underbrace{E_{\text{self}}}_{\substack{= \text{self-energy of point charges} \\ = \text{constant, but } \rightarrow \infty}}.$$

$$\Rightarrow H = \underbrace{H_{\text{matter}} + V_{\text{Coul}}}_{\equiv H'_{\text{matter}}} + H_{\text{rad}}.$$

5.2.2 Quantization

Consider again system of N point charges as before.

Qm. states of matter particles:

- Operators according to the correspondence principle:

$$\begin{aligned} H'_{\text{matter}}(\{\vec{x}_n, \vec{p}_n\}) &\rightarrow \hat{H}'_{\text{matter}}\left(\{\hat{\vec{x}}_n, \hat{\vec{p}}_n\}\right) \\ &= \sum_n \frac{1}{2m_n} \left(\hat{\vec{p}}_n - q_n \vec{A}(\hat{\vec{x}}_n, t) \right)^2 + V_{\text{Coul}}(\{\hat{\vec{x}}_n\}), \\ \text{with } [\hat{x}_{n,a}, \hat{p}_{m,b}] &= i\hbar\delta_{mn}\delta_{ab}. \end{aligned} \quad (5.43)$$

- Many-particle states $|\psi\rangle \equiv |\psi_1\rangle \cdots |\psi_n\rangle$ as direct products (anti-symmetrized/symmetrized if needed).
- Operators and states defined in the Schrödinger or Heisenberg picture as usual.

Quantized radiation field: (Coulomb gauge $\vec{\nabla} \vec{A} = 0$)

New: Φ does not vanish, because $\Delta\Phi = -\vec{\nabla} \vec{E}^\parallel = -\rho/\epsilon_0 \neq 0$.

But: Φ completely determined by charge distribution ρ , see Eq. (5.39).

↪ Φ and \vec{E}^\parallel are not dynamical variables.

Dynamical variables: $\vec{A}(\vec{x}, t)$ and $\vec{\Pi}(\vec{x}, t) = \epsilon_0 \dot{\vec{A}}(\vec{x}, t) = -\epsilon_0 \vec{E}^\perp(\vec{x}, t)$.

- Justification of $\vec{\Pi}$ as canonical mom. variable by verifying EOMs (Maxwell's eqs.).

Note: Definition formally identical to free-field case, because interactions do not involve $\dot{\vec{A}}$.

- Promotion to field operators: $\vec{A}(\vec{x}, t) \rightarrow \hat{\vec{A}}(\vec{x}, t)$, $\vec{\Pi}(\vec{x}, t) \rightarrow \hat{\vec{\Pi}}(\vec{x}, t)$.

Note: $\hat{\vec{A}}$ and $\hat{\vec{\Pi}}$ obey canonical EOMs (in the Heisenberg picture) and thus do not have simple plane-wave expansions as for the free fields given in Eq. (5.24).

- Hamiltonian:

$$\hat{H}_{\text{rad}}\left(\hat{\vec{A}}, \hat{\vec{\Pi}}\right) = \int_V d^3x \frac{1}{2} \left[\hat{\vec{\Pi}}(\vec{x}, t)^2 / \epsilon_0 + (\vec{\nabla} \times \hat{\vec{A}}(\vec{x}, t))^2 / \mu_0 \right]. \quad (5.44)$$

- Equal-time commutators:

$$\left[\hat{A}_a(\vec{x}, t), \hat{\Pi}_b(\vec{y}, t) \right] = i\hbar \delta_{ab}^\perp(\vec{x} - \vec{y}). \quad (5.45)$$

- Photon states $|\phi\rangle \in \mathcal{F}$, but states with fixed number of photons \neq eigenstates of \hat{H} .

Complete system:

- Hamiltonian:

$$\hat{H} = \hat{H}'_{\text{matter}}(\{\hat{\vec{x}}_n, \hat{\vec{p}}_n\}) + \hat{H}_{\text{rad}}(\hat{\vec{A}}, \hat{\vec{\Pi}}). \quad (5.46)$$

- Composite states:

$$|\Psi\rangle = \underbrace{|\psi\rangle}_{\text{matter part}} \otimes \underbrace{|\phi\rangle}_{\text{photon part}} = |\psi\rangle |\phi\rangle \in \mathcal{H} = \mathcal{H}_{\text{matter}} \otimes \mathcal{F}. \quad (5.47)$$

Note:

- ◊ $\hat{\vec{x}}_n, \hat{\vec{p}}_n$ act on $|\psi\rangle$,
- ◊ $\hat{\vec{A}}, \hat{\vec{\Pi}}$ act on $|\phi\rangle$.
- ◊ $[\hat{\vec{x}}_n, \hat{\vec{A}}(\hat{\vec{x}}_m, t)] = [\hat{\vec{x}}_n, \hat{\vec{\Pi}}(\hat{\vec{x}}_m, t)] = 0$.

- Interaction picture:

- ◊ Free Hamiltonian:

$$\hat{H}_0 = \sum_n \frac{\hat{\vec{p}}_n^2}{2m_n} + \hat{H}_{\text{rad}} = \text{Hamiltonian for free particles / photons.} \quad (5.48)$$

↪ Mode decomposition (5.24) of $\hat{\vec{A}}(\vec{x}, t)$ and N -photon states (5.37) correspond to exact field operators and states of the unperturbed system.

- ◊ Interaction Hamiltonian (“perturbation”):

$$\hat{H}_{\text{int}} = \sum_n \left(-\frac{q_n}{m_n} \hat{\vec{A}}(\hat{\vec{x}}_n, t) \hat{\vec{p}}_n + \frac{q_n^2}{2m_n} \hat{\vec{A}}(\hat{\vec{x}}_n, t)^2 \right) + V_{\text{Coul}}(\{\hat{\vec{x}}_n\}). \quad (5.49)$$

- Other variants of splitting \hat{H} into \hat{H}_0 and \hat{H}_{int} possible.

Example: e^- in atoms

↪ V_{Coul} considered as part of \hat{H}_0 .

5.2.3 Application: $1e^-$ atoms in quantized radiation field

Recapitulation of classical elmg. field (see end of Section III.4)

- Classical radiation field: (monochromatic, Coulomb gauge)

$$\vec{A}(\vec{x}, t) = A_0 \vec{\varepsilon} \cos(\vec{k}\vec{x} - \omega t), \quad \vec{k}\vec{\varepsilon} = 0. \quad (5.50)$$

Average energy density:

$$\bar{\rho}_E = 2\epsilon_0 \omega^2 A_0^2. \quad (5.51)$$

- Unperturbed qm. system: $1e^-$ in Coulomb field of nucleus

$$\hat{H}_0 = \frac{\hat{p}^2}{2m_e} + V_{\text{Coul}}(\hat{\vec{x}}). \quad (5.52)$$

- Perturbation: interaction with classical \vec{A} to linear order

$$\begin{aligned} \hat{H}_{\text{int}} &= \frac{e}{m_e} \vec{A}(\hat{\vec{x}}, t) \hat{\vec{p}} \\ &= \hat{h} e^{-i\omega t} + \hat{h}^\dagger e^{i\omega t}, \quad \hat{h} = \frac{e}{m_e} e^{i\vec{k}\vec{x}} A_0 \vec{\varepsilon} \hat{\vec{p}}. \end{aligned} \quad (5.53)$$

- Amplitudes for atomic transition $|i\rangle \rightarrow |f\rangle$: ($|i\rangle, |f\rangle$ = atomic energy eigenstates)

$$\begin{aligned} h_{fi} &= \langle f | \hat{h} | i \rangle = \frac{e}{m_e} A_0 \underbrace{\langle f | e^{i\vec{k}\hat{\vec{x}}} \vec{\varepsilon} \hat{\vec{p}} | i \rangle}_{\equiv M_{fi}(\vec{k})}, \\ h_{fi}^\dagger &= \langle f | \hat{h}^\dagger | i \rangle = \frac{e}{m_e} A_0 M_{fi}(-\vec{k}). \end{aligned} \quad (5.54)$$

- Rates of absorption / stimulated emission:

$$\begin{aligned} \frac{W_{fi,\text{abs}}}{T} &= \frac{2\pi}{\hbar} |h_{fi}|^2 \delta(E_f - E_i - \hbar\omega) = \frac{4\pi^2 \alpha c \bar{\rho}_E}{m_e^2 \omega^2} |M_{fi}(\vec{k})|^2 \delta(E_f - E_i - \hbar\omega), \\ \frac{W_{fi,\text{st.em}}}{T} &= \frac{2\pi}{\hbar} |h_{fi}^\dagger|^2 \delta(E_f - E_i + \hbar\omega) = \frac{4\pi^2 \alpha c \bar{\rho}_E}{m_e^2 \omega^2} |M_{fi}(-\vec{k})|^2 \delta(E_f - E_i + \hbar\omega). \end{aligned} \quad (5.55)$$

↪ Further manipulations with dipole approximation for M_{fi} , average over polarization, and integration over frequency spectrum of radiation.

Interaction with quantized radiation:

- Quantized radiation field: (single mode, Coulomb gauge)

$$\hat{\vec{A}}(\vec{x}, t) = \sqrt{\frac{\hbar}{2\omega\epsilon_0 V}} \left(\hat{a}(\vec{k}) \vec{\varepsilon}(\vec{k}) e^{i\vec{k}\vec{x} - i\omega t} + \hat{a}(\vec{k})^\dagger \vec{\varepsilon}^*(\vec{k}) e^{-i\vec{k}\vec{x} + i\omega t} \right), \quad \vec{k}\vec{\varepsilon} = 0. \quad (5.56)$$

Average energy density:

$$\bar{\rho}_E = \frac{N}{V} \hbar\omega, \quad N = \text{number of photons in volume } V. \quad (5.57)$$

- Unperturbed qm. system: $1e^-$ in Coulomb field of nucleus + free radiation field

$$\hat{H}_0 = \frac{\hat{\vec{p}}^2}{2m_e} + V_{\text{Coul}}(\hat{\vec{x}}) + \hat{H}_{\text{rad}}. \quad (5.58)$$

- Perturbation: interaction with quantized $\hat{\vec{A}}$ to linear order

$$\begin{aligned} \hat{H}_{\text{int}} &= \frac{e}{m_e} \hat{\vec{A}}(\hat{\vec{x}}, t) \hat{\vec{p}} \\ &= \hat{h} e^{-i\omega t} + \hat{h}^\dagger e^{i\omega t}, \quad \hat{h} = \frac{e}{m_e} \sqrt{\frac{\hbar}{2\omega\epsilon_0 V}} e^{i\vec{k}\hat{\vec{x}}} \hat{a}(\vec{k}) \vec{\varepsilon}(\vec{k}) \hat{\vec{p}}. \end{aligned} \quad (5.59)$$

- Amplitudes for transition $|\Psi_i\rangle \rightarrow |\Psi_f\rangle$: $|\Psi_i\rangle = |i\rangle |\phi_i\rangle$, $|\Psi_f\rangle = |f\rangle |\phi_f\rangle$,

$$\begin{aligned} h_{fi} &= \langle \Psi_f | \hat{h} | \Psi_i \rangle = \frac{e}{m_e} \sqrt{\frac{\hbar}{2\omega\epsilon_0 V}} \underbrace{\langle \phi_f | \hat{a}(\vec{k}) | \phi_i \rangle}_{\equiv M_{fi}(\vec{k})} \underbrace{\langle f | e^{i\vec{k}\vec{x}} \vec{\varepsilon} \hat{\vec{p}} | i \rangle}_{\equiv M_{fi}(\vec{k})}, \\ h_{fi}^\dagger &= \langle \Psi_f | \hat{h}^\dagger | \Psi_i \rangle = \frac{e}{m_e} \sqrt{\frac{\hbar}{2\omega\epsilon_0 V}} \langle \phi_f | \hat{a}^\dagger(\vec{k}) | \phi_i \rangle M_{fi}(-\vec{k}). \end{aligned} \quad (5.60)$$

Photon states:

$$\begin{aligned} |\phi_i\rangle &= \frac{(\hat{a}^\dagger(\vec{k}))^N}{\sqrt{N!}} |0\rangle, \quad \text{normalized } N\text{-photon state,} \\ |\phi_f\rangle &= \frac{(\hat{a}^\dagger(\vec{k}))^{N'}}{\sqrt{N'!}} |0\rangle. \end{aligned} \quad (5.61)$$

$$\Rightarrow \quad \langle \phi_f | \hat{a}(\vec{k}) | \phi_i \rangle = \underbrace{\sqrt{N} \delta_{N',N-1}}_{\text{1-photon absorption}}, \quad \langle \phi_f | \hat{a}^\dagger(\vec{k}) | \phi_i \rangle = \underbrace{\sqrt{N+1} \delta_{N',N+1}}_{\text{1-photon emission}}. \quad (5.62)$$

Squared amplitudes:

$$\begin{aligned} |h_{fi}|^2 &= \frac{\hbar e^2}{2\omega\epsilon_0 m_e^2} \frac{N}{V} |M_{fi}(\vec{k})|^2 = \frac{e^2 \bar{\rho}_E}{2\omega^2 \epsilon_0 m_e^2} |M_{fi}(\vec{k})|^2, \\ |h_{fi}^\dagger|^2 &= \frac{\hbar e^2}{2\omega\epsilon_0 m_e^2} \frac{N+1}{V} |M_{fi}(-\vec{k})|^2 \\ &= \frac{e^2 \bar{\rho}_E}{2\omega^2 \epsilon_0 m_e^2} |M_{fi}(-\vec{k})|^2 + \frac{\hbar e^2}{2\omega\epsilon_0 m_e^2} \frac{1}{V} |M_{fi}(-\vec{k})|^2. \end{aligned} \quad (5.63)$$

- Rates of absorption / stimulated emission as for classical radiation:

$$\begin{aligned} \frac{W_{fi,\text{abs}}}{T} &= \frac{4\pi^2 \alpha c \bar{\rho}_E}{m_e^2 \omega^2} |M_{fi}(\vec{k})|^2 \delta(E_f - E_i - \hbar\omega), \\ \frac{W_{fi,\text{st.em}}}{T} &= \frac{4\pi^2 \alpha c \bar{\rho}_E}{m_e^2 \omega^2} |M_{fi}(-\vec{k})|^2 \delta(E_f - E_i + \hbar\omega). \end{aligned} \quad (5.64)$$

NEW: contribution for “spontaneous emission” (independent of $\bar{\rho}_E$!):

$$\frac{W_{fi,\text{sp.em}}}{T} = \frac{\pi e^2}{\epsilon_0 \omega m_e^2} \frac{1}{V} |M_{fi}(-\vec{k})|^2 \delta(E_f - E_i + \hbar\omega) \quad (5.65)$$

= transition rate for emitting a photon
with specific momentum $\hbar k$ and polarization $\vec{\varepsilon}$

(5.66)

- Decay width $\Gamma_{fi} = \hbar \times \text{rate}$ for the atomic transition $|i\rangle \rightarrow |f\rangle$ via spontaneous emission of any photon in dipole approximation:

$$\begin{aligned} \Gamma_{fi} &= \sum_{\vec{k}} \sum_{\gamma \text{ pol}} \frac{\pi \hbar e^2}{\epsilon_0 \omega m_e^2} \frac{1}{V} |M_{fi}(-\vec{k})|^2 \delta(E_f - E_i + \hbar\omega) \\ &= \underbrace{\frac{1}{V} \sum_{\vec{k}}}_{\int \frac{d^3k}{(2\pi)^3}} \frac{\pi \hbar e^2}{\epsilon_0 \omega m_e^2} \underbrace{\sum_{\gamma \text{ pol}} |M_{fi}(-\vec{k})|^2}_{= 2m_e^2 \omega_{if}^2 \vec{x}_{fi}^2 / 3 \text{ in dipole approximation},} \delta(E_f - E_i + \hbar\omega) \\ &= \int \frac{d^3k}{(2\pi)^3} \frac{2\pi e^2 \omega_{if}^2 \vec{x}_{fi}^2}{3\epsilon_0 \omega} \delta(\omega - \omega_{if}), \quad \omega = ck \\ &= \frac{e^2 \omega_{if}^3 \vec{x}_{fi}^2}{3\pi \epsilon_0 c^3} = \frac{4\alpha \hbar \omega_{if}^3 \vec{x}_{fi}^2}{3c^2}. \end{aligned} \quad (5.67)$$

Total decay width (“natural width”) Γ_i of atomic state $|i\rangle$:

$$\Gamma_i = \frac{\hbar}{\tau_i} = \sum_{\substack{f \\ E_f < E_i}} \Gamma_{fi}, \quad \tau_i = \text{lifetime of } |i\rangle. \quad (5.68)$$