

Exercise 5.1 (1 point) *Integration over Grassmann variables*

We consider a Grassmann algebra with the $2N$ generating elements y_k, y_k^* ($k = 1, \dots, N$).

- a) The generators y_k will be replaced by new generators z_k with the mapping $z_i = a_{ik}y_k$, where a_{ik} denote the coefficients of a complex matrix A . Show that the corresponding differentials transform as $dz_i = \tilde{a}_{ik}dy_k$, where \tilde{a}_{ik} are the coefficients of the matrix that is the transposed inverse matrix of A .
- b) Using a), derive

$$\int dz_1 \cdots \int dz_n F(\mathbf{z}) = (\det A)^{-1} \int dy_1 \cdots \int dy_n F(\mathbf{z}(\mathbf{y})).$$

- c) With the help of b) show that the Gaussian integral is given by

$$\int dy_1 \cdots \int dy_n \int dy_n^* \cdots \int dy_1^* \exp\{y_i^* a_{ik} y_k\} = \det A.$$

Exercise 5.2 (1.5 points) *Generating functional of free Dirac fields*

The generating functional of free Dirac fields $\psi, \bar{\psi}$ is given by the functional integral

$$Z_{\psi,0}[\eta, \bar{\eta}] = N \int \mathcal{D}\psi \int \mathcal{D}\bar{\psi} \exp \left\{ i \int d^4x \left[\bar{\psi}(x) (i\cancel{\partial} - m) \psi(x) + \bar{\eta}(x)\psi(x) + \bar{\psi}(x)\eta(x) \right] \right\}$$

with the normalization $Z_{\psi,0}[0, 0] = 1$. The source fields $\eta, \bar{\eta}$ are of Grassmann-type.

- a) Analogous to the procedure for scalar fields, calculate the functional $Z_{\psi,0}[\eta, \bar{\eta}]$ as

$$Z_{\psi,0}[\eta, \bar{\eta}] = \exp \left\{ \int d^4x \int d^4x' i\bar{\eta}(x) iS_F(x-x') i\eta(x') \right\},$$

where $iS_F(x)$ is defined by $(i\cancel{\partial} - m) S_F(x) = \delta(x)$.

- b) Starting from the generating functional $Z_{\psi,0}[\eta, \bar{\eta}]$, derive the fermion propagator

$$G_0^{\bar{\psi}\psi}(x_1, x_2) = \frac{\delta}{i\delta\eta(x_2)} \frac{\delta}{i\delta\bar{\eta}(x_1)} Z_{\psi,0}[\eta, \bar{\eta}] \Big|_{\eta, \bar{\eta}=0}.$$

- c) Solve the differential equation for $S_F(x)$ in momentum space and derive an explicit form of the Fourier representation of the propagator

$$G_0^{\bar{\psi}\psi}(x_1, x_2) = \int \frac{d^4q}{(2\pi)^4} e^{-iq(x_1-x_2)} \tilde{G}_0^{\bar{\psi}\psi}(q).$$

Please turn over !

Exercise 5.3 (1 point) *Green's function for the elementary quark–gluon interaction*

The generating functional of QCD can be written as

$$Z[J_\mu^a, \eta, \bar{\eta}] = N \exp \left\{ i \int d^4 y \mathcal{L}_I \left(A_\mu^a(y) \rightarrow \frac{\delta}{i \delta J^{\mu,a}(y)}, \bar{\psi}(y) \rightarrow -\frac{\delta}{i \delta \eta(y)}, \psi(y) \rightarrow \frac{\delta}{i \delta \bar{\eta}(y)} \right) \right\} \\ \times Z_{A,0}[J_\mu^a] Z_{\psi,0}[\eta, \bar{\eta}],$$

$$Z[0, 0, 0] = 1,$$

$$Z_{A,0}[J_\mu^a] = \exp \left\{ \frac{1}{2} \int d^4 x \int d^4 x' i J^{\mu,a}(x) G_{0,\mu\nu}^{AA,ab}(x, x') i J^{\nu,b}(x') \right\},$$

$$Z_{\psi,0}[\eta, \bar{\eta}] = \exp \left\{ \int d^4 x \int d^4 x' i \bar{\eta}(x) G_0^{\bar{\psi}\psi}(x, x') i \eta(x') \right\},$$

where the Lagrangian for the quark–gluon interaction is given by $\mathcal{L}_I = -g \bar{\psi} A_\mu^a T^a \gamma^\mu \psi$. Calculate the Green's function

$$G_\mu^{A\bar{\psi}\psi,a}(x_1, x_2, x_3) = \frac{\delta}{i \delta J^{\mu,a}(x_1)} \frac{\delta}{i \delta \eta(x_3)} \frac{\delta}{i \delta \bar{\eta}(x_2)} Z[J_\mu^a, \eta, \bar{\eta}] \Big|_{J_\mu^a, \eta, \bar{\eta} \rightarrow 0}$$

in lowest-order perturbation theory and, after transforming into momentum space, verify that the corresponding amputated Green's function agrees with the Feynman rule $-ig T^a \gamma_\mu$.