
Exercises to Relativistic Quantum Field Theory — Sheet 1

— Prof. S. Dittmaier, Universität Freiburg, SS18 —

Exercise 1.1 *Some properties of Lorentz transformations* (2 points)

Lorentz transformations, which transform a four-vector $a^\mu = (a^0, \vec{a})$ to $a'^\mu = \Lambda^\mu{}_\nu a^\nu$, comprise all 4×4 matrices Λ that leave the metric tensor $g^{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ invariant, i.e. $g^{\mu\nu} = \Lambda^\mu{}_\alpha \Lambda^\nu{}_\beta g^{\alpha\beta}$. In the following we consider the “proper orthochronous Lorentz group” L_+^\uparrow that comprises all such Λ with the two constraints that $\det \Lambda = +1$ and $\Lambda^0{}_0 > 0$. The group L_+^\uparrow consists of all rotations in space and “boosts”, which relate two frames of reference with a non-vanishing relative velocity.

- a) A boost with relative velocity $\vec{v} = (v^1, v^2, v^3)$ is described by the Λ matrix of the form

$$L(v^1, v^2, v^3) = \begin{pmatrix} \gamma & -v^1\gamma & -v^2\gamma & -v^3\gamma \\ -v^1\gamma & 1 + \frac{v^1 v^1}{\vec{v}^2}(\gamma - 1) & \frac{v^1 v^2}{\vec{v}^2}(\gamma - 1) & \frac{v^1 v^3}{\vec{v}^2}(\gamma - 1) \\ -v^2\gamma & \frac{v^2 v^1}{\vec{v}^2}(\gamma - 1) & 1 + \frac{v^2 v^2}{\vec{v}^2}(\gamma - 1) & \frac{v^2 v^3}{\vec{v}^2}(\gamma - 1) \\ -v^3\gamma & \frac{v^3 v^1}{\vec{v}^2}(\gamma - 1) & \frac{v^3 v^2}{\vec{v}^2}(\gamma - 1) & 1 + \frac{v^3 v^3}{\vec{v}^2}(\gamma - 1) \end{pmatrix},$$

where $\gamma = 1/\sqrt{1 - \vec{v}^2}$. Calculate the boosted components x'^μ for the four-vectors $x_{\parallel} = (x^0, r\vec{e})$ and $x_{\perp} = (x^0, r\vec{e}_{\perp})$ whose directions in space are parallel and perpendicular to the direction $\vec{e} = \vec{v}/|\vec{v}|$ of the relative velocity, respectively, i.e. $\vec{e}_{\perp} \cdot \vec{e} = 0$.

- b) Show that the sign of the time-like component a^0 of any non-space-like four-vector a^μ (i.e. $a^2 \geq 0$) is invariant under all Lorentz transformations $\Lambda \in L_+^\uparrow$.
- c) Calculate $W = L_2(0, -v^2, 0)L_1(-v^1, 0, 0)L_2(0, v^2, 0)L_1(v^1, 0, 0)$ for small velocities v^k and keep terms up to quadratic order in products of components v^k . What kind of transformation is described by W ?
- d) Show that the totally antisymmetric tensor

$$\epsilon^{\mu\nu\rho\sigma} = \begin{cases} +1 & \text{if } (\mu\nu\rho\sigma) = \text{even permutation of } (0123), \\ -1 & \text{if } (\mu\nu\rho\sigma) = \text{odd permutation of } (0123), \\ 0 & \text{otherwise.} \end{cases}$$

is an invariant tensor under all $\Lambda \in L_+^\uparrow$, i.e. $\epsilon'^{\mu\nu\rho\sigma} = \Lambda^\mu{}_\alpha \Lambda^\nu{}_\beta \Lambda^\rho{}_\gamma \Lambda^\sigma{}_\delta \epsilon^{\alpha\beta\gamma\delta}$.

- e) Show that $d^\mu = \epsilon^{\mu\nu\rho\sigma} a_\nu b_\rho c_\sigma$ transforms like a four-vector under $\Lambda \in L_+^\uparrow$ if a^μ , b^μ , and c^μ are four-vectors.

Please turn over!

Exercise 1.2 *Kinematics of a $1 \rightarrow 2$ particle decay* (2 points)

A particle of mass M and four-momentum k^μ decays into two particles of masses m_i and four-momenta p_i^μ ($i = 1, 2$). The momenta obey their mass-shell conditions $k^2 = M^2$ and $p_i^2 = m_i^2$ and, in the centre-of-mass frame Σ , are given by

$$k^\mu = (M, \vec{0}), \quad p_i^\mu = (E_i, |\vec{p}_i| \cos \phi_i \sin \theta_i, |\vec{p}_i| \sin \phi_i \sin \theta_i, |\vec{p}_i| \cos \theta_i).$$

- a) What are the consequences of four-momentum conservation $k = p_1 + p_2$ for the energies E_i , for the absolute values $|\vec{p}_i|$ of the three-momenta and for the angles θ_i , ϕ_i ?
- b) Calculate E_i and $|\vec{p}_i|$ as functions of the masses M and m_i .
- c) The decaying particle is now considered in a frame Σ' in which the particle has the velocity β along the x^3 axis ($c = 1$). What is the relation between energies and angles in Σ' with the respective quantities in Σ ?
- d) For the special case $m_1 = m_2 = 0$ (e.g. decay into two photons) determine the angle θ' between the directions of flight of the decay products in Σ' (i.e. the angle between \vec{p}'_1 and \vec{p}'_2). What are the extremal values of θ' ? In particular, discuss the cases $\beta = 0$ and $\beta \rightarrow 1$.